

An Overview of the 13:8 Mean Motion Resonance between Venus and Earth

Á. Bazsó · V. Eybl · R. Dvorak · E. Pilat-Lohinger · Ch. Lhotka

Received: date / Accepted: date

Abstract It is known since the seminal study of Laskar (1989) that the inner planetary system is chaotic with respect to its orbits and even escapes are not impossible, although in time scales of billions of years. The aim of this investigation is to locate the orbits of Venus and Earth in phase space, respectively to see how close their orbits are to chaotic motion which would lead to unstable orbits for the inner planets on much shorter time scales. Therefore we did numerical experiments in different dynamical models with different initial conditions – on one hand the couple Venus-Earth was set close to different mean motion resonances (MMR), and on the other hand Venus’ orbital eccentricity (or inclination) was set to values as large as $e = 0.36$ ($i = 40^\circ$). The couple Venus-Earth is almost exactly in the 13:8 mean motion resonance. The stronger acting 8:5 MMR inside, and the 5:3 MMR outside the 13:8 resonance are within a small shift in the Earth’s semimajor axis (only 1.5 percent). Especially Mercury is strongly affected by relatively small changes in eccentricity and/or inclination of Venus in these resonances. Even escapes for the innermost planet are possible which may happen quite rapidly.

Keywords planetary motion · mean motion resonances Venus and Earth · Mercury’s escape

1 Introduction

Mean Motion Resonances are essential in Solar System Dynamics not only for the planets but also for the motion of the asteroids. The appearance of chaotic motion in the asteroid belt detected by Wisdom (1981) – the 3:1 MMR of an asteroid with Jupiter – was the first one discovered after the seminal discovery by Henon and Heiles (1964) for galactic dynamics. But the structure of the asteroid belt is also created by the secular resonances (SR), where the motion of the perihelia and the nodes can be in resonance for different planets in combination with the asteroid’s perihelion respectively node. For the planets it was found by Laskar in different papers (Laskar (1988, 1990, 1996)) that especially the Inner Solar System (ISS) is in a chaotic state. The appearance of SR may even shift the orbit of Mars out from a stable orbit into an unstable one with high probability when one does not take into account relativity. The large planets seem to be in quite a ‘safe’ region, although




Fig. 1 The coupling in the eccentricities of Venus and Earth; from -100 to 95 Myrs (upper graph), from -2.5 to 2.5 Myrs (middle graph) and from 95 to 100 Myrs (lower graph), after Dvorak and Gamsjäger (2003).

many MMR are acting: e.g. the 5:2 MMR between Jupiter and Saturn. Recent investigations by Michtchenko and Ferraz-Mello (2001); Gallardo (2006) show how close their motion is to a chaotic state with severe implications for the other planets.

Because the orbits of Venus and Earth are quite close to the 13:8 MMR¹ and they are strongly coupled with respect to the inclinations and eccentricities (Fig. 1) – comparable to Jupiter and Saturn – we investigated the phase space close to this high order resonance. The order of a resonance is defined as the difference q when we characterize a MMR by $p : (p + q)$. The value q is connected to the exponent in front of the Fourier term in the expansion of the disturbing function; the higher the exponent in a small quantity in front of the Fourier term is, the less strong is the influence on the dynamics, unless there is a resonance acting (e.g. Murray and Dermott). Thus for the 13:8 MMR we have to deal with the order 5. It is interesting to note that a relatively small shift of the Earth to a smaller semimajor axis would bring the couple Venus–Earth into the 8:5 MMR (order 3); a shift to a larger semimajor axis would bring both planets into the 5:3 MMR (order 2) (Fig. 2). In this plot the results of the computations in the simple three body problem Sun–Venus–Earth are presented to show the location of the resonances and also their shape.

Besides extensive numerical integration of the motions of the planets in different dynamical models, we also used chaos indicators and analysed the frequencies involved. Furthermore simplified mapping models were constructed to understand the structure of phase space where Venus and Earth are embedded.

The content of the paper is follows: after a careful test of the dynamical model which we used (section 2) we roughly describe the three MMR mentioned above and show the structure determined with the aid of a chaos indicator (section 3). We then report on the results of different numerical experiments where we show the dependency of the stability of

¹ the periods of Venus and Earth are almost in the ratio 13:8




Fig. 2 The MMRs between Venus and the Earth; along the x-axis the initial semimajor axis of the Earth is changed, along the y-axis the initial values for the eccentricity of Venus are varied. The greytone indicates the maximum eccentricity, the vertical white lines show the position of the MMR up to high orders.

the inner planetary system on the eccentricities and the inclinations of Venus (section 4). In the conclusions we discuss the possible consequences for the stability of the inner planetary system. In an appendix first results of an applied mapping for motions in MMRs are shown.

2 The dynamical models and the numerical setup

To choose a quite realistic model we first did integrations with the actual initial conditions of Venus and Earth in different dynamical models, using a numerical integration scheme based on Lie-series (Hanslmeier and Dvorak (1984); Delva (1984); Lichtenegger (1984)):

- A** the Inner Solar System (ISS, 5-bodies)
- B** the ISS + Jupiter (6-bodies)
- C** the ISS + Jupiter + Saturn (7-bodies)
- D** the ISS + Jupiter + Saturn + Uranus (8-bodies)
- E** the ISS + outer Solar system (9-bodies)

The different behaviour between the models consisting only of the inner planets (**A**) and the one with Jupiter (**B**) is visible in Fig. 3: the very regular variation limited by a maximum value of $e = 0.03$ (upper graph) – is completely destroyed by the perturbation of Jupiter (Fig. 3: lower graph). One can see three distinct sections: (i) sometimes the eccentricities almost do not change for up to 10^5 years (e.g. between 2.25 and 2.35×10^6 years for Venus and 2.55 and 2.65×10^6 years for the Earth); (ii) sometimes the changes are in antiphase (like in the upper graph of Fig. 3); and (iii) sometimes they are in phase.




Fig. 3 Coupling in the eccentricities of Venus and Earth for the model A (upper graph) and in the model B for 3 million years.




Fig. 4 Coupling in the eccentricities of Venus and Earth for the model C (upper graph) and in the model E for 3 million years.




Fig. 5 Comparison of the 5 dominating frequencies in the dynamics of Venus and Earth for the five different models **A** to **E**. On the x-axis the number of bodies is given (5 to 9), on the y-axis the frequencies (in arcsec per year) and in z-direction the amplitude is given. Always on the left is the frequency of Earth, on the right the frequency of Venus. The numbers on the x-axis show the number of bodies in the respective models; e.g. '5' means that the 5-body problem was used for the numerical integration.

Including now Saturn as perturber in model **C** changes the picture a lot: the development of the eccentricities is always in antiphase, reaches even larger maximum values, but looks 'smoother' for both planets compared to model **B** (Fig. 4 upper graph). Now by comparing the dynamical evolution of Venus and Earth including all planets of the Solar System (model **E**, Fig. 4 bottom) with **C** it is interesting to note that for the first 10^6 years even quantitatively the behaviour is very similar. Later (up to 3×10^6 years) at least qualitatively both models show an analogous behaviour².

In addition to the former computations we compare for the five different models the dominating frequencies, which were derived by the program SIGSPEC (Reegen (2007)³). SIGSPEC incorporates the frequency and phase of the signal appropriately and takes into account the properties of the time-domain sampling which provides an unprecedentedly accurate peak detection (Kallinger et al (2008)). Although the present investigations deal with equidistantly sampled time series, the benefit of frequency- and phase-resolved statistics and the capability of full automation are indispensable also in this application.

The data used were the orbital element $k = e \cos \varpi$ of the Earth and of Venus in all mentioned dynamical models. It is evident that the model **A** gives quite a wrong picture of the involved resonances. To include only Jupiter is already a much better model and taking into account the great inequality between Jupiter and Saturn **C** (x-axes '7') is a very good

² Model **D**, not shown here, is not significantly better than **C**.

³ This code is a new method in time series analysis and it is the first technique to rely on an analytic solution for the probability distribution produced by white noise in an amplitude spectrum (Discrete Fourier Transform). Returning the probability that an amplitude level is due to a random process, it provides an objective and unbiased estimator for the significance of a detected signal.




Fig. 6 Results of integrations for 10^6 years with fixed orbital elements in the 13:8 MMR: the differences in maximum eccentricity of each individual planet in 2 models (A, dashed lines, C full lines) are plotted (y-axis) for initially different inclinations of Venus (x-axis). Light grey: Earth, grey: Mars, dark grey: Mercury.

description of the motions of Earth and Venus. The differences between the models C and E are not significant as we also noticed from the direct inspection of the eccentricity plots. Consequently as compromise between long CPU time and precision of the model it is clear that C is a good choice.

One additional test was made by checking the maximum eccentricities of the orbits (e_{max}) of the inner planets when changing the initial eccentricity of Venus. The respective integrations for 10^6 years shown in Fig. 6 point out the differences in e_{max} of each individual planet in the 5 and 7-body models (A and C) for the 13:8 MMR. In contrast to the other planets one can see that the Earth (light grey) has almost always a higher maximum eccentricity due to the influence of Jupiter and Saturn than without these two planets. For Mercury and Mars no clear dependence is observable, the maximum values of their eccentricities change with the initial inclination of Venus. It is remarkable that Mercury achieves very high eccentricities, and eventually escapes for longer integrations (see next chapter).

3 Description and analysis of the 8:5, 13:8 and 5:3 MMR

In Fig. 2 we show a detailed graph of the neighborhood of the 13:8 MMR in the planar 3-body problem Sun-Venus-Earth: in addition to the 13:8, the 8:5 and the 5:3 MMR many high order MMRs are visible. To derive a simple picture of the phase space structure close to the 13:8 MMR only two initial parameters were changed: on the x-axis the semimajor axis of the Earth was varied and on the y-axis the initial eccentricity of the perturbing planet Venus is plotted. In this graph, where the maximum eccentricity of an Earth-orbit during the integration time is shown on a grey scale, one already can see the inequality of the perturbations acting in the resonances.

- The 8:5 MMR (order 3, $a = 0.989501$ AU)⁴ has a relatively narrow triangular structure with highly perturbed wings down to $e_{\text{Venus}} = 0.12$ on both sides, whereas in the middle of the resonance the orbits are very regular with small eccentricities.
- The 13:8 MMR (order 5, at $a = 0.999782$ AU) is the weakest of the three MMR we are studying. The wings on both sides of the resonance go down to $e_{\text{Venus}} = 0.15$. Again in the middle of the resonance – here broader than in the 8:5 resonance – the orbits are only marginally perturbed; the eccentricities stay well below 0.075⁵.
- The 5:3 MMR (order 2, $a = 1.016799$ AU) is by far the strongest one: it is quite broad from large eccentricities of Venus on down to small eccentricities like $e = 0.05$. It shows a big ‘quiet’ central region and in the wings the Earth’s orbit suffers from relatively large eccentricities $e_{\text{Earth}} \sim 0.3$

Another high order MMR, the 18:11 resonance, is visible close to the 13:8 resonance, which we will not discuss here because its action is evidently weaker (order 7).

To use the results of direct numerical integrations one has to carry out orbital computations over a very long time. In order to save computation time it is advisable to use a so-called chaos indicator, that shows the state of motion quite fast and allows us to reduce the integration time significantly. Therefore, we used the FAST LYAPUNOV INDICATOR (FLI) (see Froeschlé et al (1997)), which is a quite fast tool to distinguish between regular and chaotic motion. According to the definition – where Ψ is the length of the largest tangent vector:

$$\psi(t) = \sup_i \|v_i(t)\| \quad i = 1, \dots, n$$

(n denotes the dimension of the phase space) – it is obvious that chaotic orbits can be found very quickly because of the exponential growth of this vector in the chaotic region. This method has often been applied to studies of Extra-solar Planetary Systems (Lohinger et al (2008); Schwarz et al (2009)). In this study the FLIs were computed for about 50000 yrs. The resulting stability maps are shown in Figs. 7 and 8, where the black region marks the stable motion and grey areas show chaotic regions. Fig. 7 shows the state of motion from 0.98 AU to 1.020 AU for different eccentricities of the Earth. One can clearly see perturbations in the stable region even for low eccentric motion of the Earth due to MMR with Venus. The three MMRs – 8:5, 13:8 and 5:3 – that are analyzed in this study are clearly visible in Fig. 7. A magnification of the 13:8 MMR is shown in Fig. 8, which indicates chaotic behaviour for the position of the Earth if $e_{\text{Earth}} > 0.1$. The interesting fine structures inside the MMR (symmetric with respect to the central line of the resonance) unveil a central region for $e = 0.2$ which still shows regular motions of the Earth, while close to it chaotic motion is present (white in Fig. 7).

4 Dependence on the initial conditions of Venus

To check the sensitivity with respect to different initial conditions of the ‘planet twins’ Venus–Earth concerning the stability of the inner Solar System we performed a series of numerical integrations in the model C for 10^7 years where we checked the three MMR described before. Two cuts through the exact location of the MMR have been undertaken, one

⁴ with $a_{\text{Venus}} = 0.723330$ AU

⁵ The Earth with a semimajor axis $a = 1$ is just a little outside of this MMR.




Fig. 7 FLI map: semimajor axes of the Earth versus different eccentricities of Venus. Visible from left to right the 8:5, the 13:8 and the 5:3 MMR. Black regions indicate stable orbits.

where we changed the eccentricity of Venus and another one where we changed its inclination. Both results were analysed through the simple test of the largest eccentricity of the 4 inner planets during the integration.

4.1 Dependence of the Earth's orbit on the eccentricity of Venus

- **The 8:5 MMR** (Fig. 9, upper graph):

The orbits of Venus and the Earth develop qualitatively in the same way, and no dramatic changes are visible in their orbits. Mercury is the planet which suffers – even for relatively small initial eccentricities of Venus ($e_{\text{Venus}} \leq 0.12$) – from large eccentricities. For all larger values of e_{Venus} Mercury is in quite a chaotic orbit. Mars remains longer with orbital elements which do not allow close encounters to the Earth, but from $e_{\text{Venus}} \geq 0.2$ on its orbit is also destabilized.

- **13:8 MMR** (Fig. 9, middle graph):

Up to $e_{\text{Venus}} \leq 0.15$ the inner planetary system seems to be in a relatively 'quiet' region of phase space; from that on, Mercury and Mars getting larger and larger eccentricities, the system is destabilized. Venus and Earth are still developing in quite a similar way without dramatic changes of their orbits.

- **The 5:3 MMR** (Fig. 9, lower graph):

The orbit of the Earth suffers more and more from larger and larger eccentricities depending on the one of Venus very similar to the other two resonances. The two planets




Fig. 8 Zoom of the 13:8 MMR, captions like in Fig. 7.

Mercury and Mars achieve such large eccentricities that encounters with Venus respectively Earth cause them to leave their orbits. The chaotic behaviour of the orbits of the inner planetary system is already visible in the FLI diagram for $e_{\text{Venus}} \geq 0.17$ on (compare Fig. 7).

The dynamical behaviour of orbits inside these three resonances is quite similar; although the 5:3 and 8:3 MMR are of a lower order compared to the 13:8 MMR and one expects stronger perturbations on the orbits of the planets; in the respective plots this is not visible.

4.2 Dependence of the Earth's orbit on the inclination of Venus

– **The 8:5 MMR** (Fig. 10, upper graph):

In this resonance it is evident that Mercury is thrown out from its orbit already for moderate initial inclinations of Venus, although a small window of more stable orbits appears for $11^\circ \leq i \leq 12^\circ$. Earth and Venus are strongly perturbed: from $i = 29^\circ$ on eccentricities up to $e = 0.35$ for Venus and $e = 0.3$ for the Earth are possible. Mars obviously is not so much affected by the Earth, only for $i = 37^\circ$ a maximum value of $e = 0.4$ is reached, which brings him into an Earth-crossing orbit. For longer integration time again an escape seems possible. Then, surprisingly, Mars for $i = 38^\circ$ and $i = 39^\circ$ (within the integration time of 10^7 years!) still stays in a moderately elliptic orbit.

– **The 13:8 MMR** (Fig. 10, middle graph):




Fig. 9 e_{max} cuts through the centers of the 8:5, 13:8 and 5:3 MMR (from top to bottom) depending on the eccentricity of the perturbing planet Venus (x-axis).

It is quite remarkable that even a small shift of the initial inclination to $i \sim 8^\circ$ of Venus may lead to an escape of Mercury (after $e = 0.75$ a subsequent escape from its orbit because of encounters with Venus is expected). A second 'dangerous' inclination is visible around $i = 15^\circ$; then, from $i \geq 20^\circ$ on, Mercury would suffer from close approaches to Venus. There seems to be little influence on the orbits of Venus, Earth and Mars up to $i_{Venus} = 35^\circ$.

– **The 5:3 MMR** (Fig. 10, lower graph):




Fig. 10 $e_{\max}$ cuts through the centers of the 8:5, 13:8 and 5:3 MMR (from top to bottom) depending on the inclination of the perturbing planet Venus (x-axis).

The strongest of the three MMR shows in fact a dynamical evolution which is dominated by the very strong perturbations on Mercury for inclinations of Venus around 10° and then, from 15° on the large eccentricities of Mercury ($e \geq 0.6$) will always lead to escapes of this planet. Mars is not affected dramatically in this resonance, Venus and Earth stay within moderate eccentricities for all initial inclinations of Venus.

5 Conclusions

In this investigation we explored the environment of the 13:8 MMR, which is very close to the actual position of Venus and Earth. After a careful test of a simplified dynamical model of our planetary system appropriate for our task we discussed the main structures of the three MMR in the vicinity of the two planets using the results of a study of the FLI.

In another step, using extensive numerical integrations we changed the ratio of the semi-major axes Venus–Earth⁶; the orbit of the Earth was given different values of a to cover the proximate neighborhood of the MMR where the twin-planets are in. Then, in two different runs the eccentricity of Venus was set to eccentricities up to $e = 0.36$ and in the dynamical model, consisting of the inner Solar System with Jupiter and Saturn, the equations of motion were integrated for 10^7 years. The same was done in a second run where we changed the inclination of Venus up to $i = 40^\circ$. These two experiments were undertaken for each of the three MMR 8:5, 13:8 and 5:3 and the results analysed with respect to the maximum eccentricities of the inner planets. It turned out that only Mercury is the planet which would suffer from such perturbations, that it would achieve very large values of its eccentricity and thus would be thrown out from its current orbit. According to our results Mars is not as strongly affected by a larger inclination or eccentricity of Venus as Mercury; this can be understood by the presence of Jupiter and Saturn keeping this planet almost in its original plane of motion. Venus and also the Earth are only affected for larger initial inclinations of Venus, but the planets stay more or less in their orbits although sometimes achieving eccentricities as large as $e = 0.3$.

In an attempt to understand the dynamical structure of the resonances a mapping approach (appendix) was constructed in the simple elliptic restricted three body problem Sun–Venus and massless Earth for the 3:1 MMR. The same approach was then used to construct the surface of section of the 5:3 MMR. Until now we failed to get a similar graph for the other two resonances which ask for a development of the Hamiltonian up to order 3 and order 5, but we work on it.

As a final statement we may say that according to our results although the couple of Venus and Earth is close to a MMR we do not expect big changes of the orbits of the inner planets unless Venus will come into an orbit of an inclination of about 7 degrees or will achieve an eccentricity of 0.2. Then, in fact, Mercury would be on an unstable orbit which could finally throw it either into the Sun or far out into the main belt of asteroids!

6 Appendix: A Mapping Model

Mapping methods are very efficient analytical tools for studying the dynamics of mean motion resonances. Their main advantage is the fact, that a mapping is much faster in regard to computing time, while also being more exact than numerical integrations in the same model. In this work, we followed the approach used in the works of Hadjidemetriou (1993); Tsiganis (2007); Lhotka et al (2008), which has been found very successful already in the study of the 1:1 resonance. The mapping allows us to investigate the phase space structures of $p : (p + q)$ MMR in a semi-analytical way. The Hamiltonian of the three body problem in the exact location of the chosen resonance is used to derive equations, which provide a set of rules for the propagation of a trajectory in phase space. The Hamiltonian in Keplerian elements (a, e, ω, Ω) describing the motion of the perturbed body is averaged over the

⁶ Venus' semimajor axis was fixed.




Fig. 11 Phase space portrait of the 3:1 resonance (left) and 5:3 resonance (right) of a massless body with Earth.

orbital period of the perturbing body and then transformed to action and angle variables (X, Y, ψ, J) in phase space by canonical transformations. The mapping equations then provide iteration rules for each of these variables with a time step equaling the orbital period of the perturber. Being based on the exact equations of motion, the results obtained from the mapping equations provide the most accurate insight in the dynamics of the chosen system.

To test the applicability of the method, the 3:1 resonance of a massless body with Jupiter in the planar elliptic restricted three body problem has been analyzed, which yielded very good results. The same method was used for the 5:3 resonance of a massless particle with Earth, again in the planar ER3BP. These two resonances are of order two and therefore more easily computable than the 8:5 and the 13:8 resonance. In Fig. 11 (left graph) the phase space portrait for the 3:1 mean motion resonance of a test body with Jupiter is shown. The same method was applied to the 5:3 MMR with Earth. Up to now, we have been studying only low-order resonances in the planar elliptic restricted three body problem using the described method with good results. In a next step we will expand this work to the investigation of the 8:5 MMR and especially the 13:8 MMR, which was the task of this study.

7 Acknowledgements

A. Bazso, V. Eybl and Ch. Lhotka appreciate the financial help of the FWF project P 18930-N16; E. Pilat-Lohinger needs to thank the FWF project P-19569-N16.

References

- Delva, M., "Integration of the elliptic restricted three-body problem with Lie series", *Celest. Mech.*, 34, 145-154, (1984)
- Dvorak, R. and Süli, Á., "On the Stability of the Terrestrial Planets as Models for Exosolar Planetary Systems", *Celest. Mech. Dyn. Astron.*, 83, 77-95, (2002)
- Dvorak, R. and Gamsjäger, C., "A New Determination of the Basic Frequencies in Planetary Motion", *Proceedings of the 3rd Austrian-Hungarian Workshop on Trojans and related Topics*, 49-58. F. Freistetter, R. Dvorak and B. Érdi (Eds.), (2003)
- Froeschlé, C. and Lega, E. and Gonczi, R., "Fast Lyapunov Indicators. Application to Asteroidal Motion", *Celest. Mech. Dyn. Astron.*, 67, 41-62, (1997)
- Gallardo, T., "Atlas of the mean motion resonances in the Solar System", *Icarus*, 184, 29-38, (2006)
- Hadjimetriou, J. D., "Asteroid motion near the 3:1 resonance", *Celest. Mech. Dyn. Astron.*, 56, 563-599, (1993)
- Hansmeier, A. and Dvorak, R., "Numerical Integration with Lie Series", *A&A*, 132, 203-+, (1984)

- Henon, M. and Heiles, C., "The applicability of the third integral of motion: Some numerical experiments", *AJ*, 69, 73-+, (1964)
- Kallinger, T. and Reegen, P. and Weiss, W. W., "A heuristic derivation of the uncertainty for frequency determination in time series data", *A&A*, 481, 571-574, (2008)
- Laskar, J., "Secular evolution of the solar system over 10 million years", *A&A*, 198, 341-362, (1988)
- Laskar, J., "A numerical experiment on the chaotic behaviour of the solar system", *Nature*, 338, 237-+, (1989)
- Laskar, J., "The chaotic motion of the solar system - A numerical estimate of the size of the chaotic zones", *Icarus*, 88, 266-291, (1990)
- Laskar, J., "Large-scale chaos in the solar system", *A&A*, 287, L9-L12, (1994)
- Laskar, J., "Large Scale Chaos and Marginal Stability in the Solar System", *Celest. Mech. Dynam. Astron.*, 64, 115-162, (1996)
- Laskar, J., "Large scale chaos and the spacing of the inner planets.", *A&A*, 317, L75-L78, (1997)
- Lhotka, C. and Efthymiopoulos, C. and Dvorak, R., "Nekhoroshev stability at L_4 or L_5 in the elliptic-restricted three-body problem - application to Trojan asteroids", *MNRAS*, 384, 1165-1177, (2008)
- Lichtenegger, H., "The dynamics of bodies with variable masses", *Celest. Mech.*, 34, 357-368, (1984)
- Pilat-Lohinger, E. and Robutel, P. and Süli, Á. and Freistetter, F., "On the stability of Earth-like planets in multi-planet systems", *Celest. Mech. Dyn. Astron.*, 102, 83-95, (2008)
- Michtchenko, T. A. and Ferraz-Mello, S., "Modeling the 5 : 2 Mean-Motion Resonance in the Jupiter-Saturn Planetary System", *Icarus*, 149, 357-374, (2001)
- Reegen, P., "SigSpec. I. Frequency- and phase-resolved significance in Fourier space", *A&A*, 467, 1353-1371, (2007)
- Schwarz, R. and Süli, Á. and Dvorak, R. and Pilat-Lohinger, E., "Stability of Trojan planets in multi-planetary systems. Stability of Trojan planets in different dynamical systems", 104, 69-84, (2009)
- Tsiganis, K., "Chaotic Diffusion of Asteroids", *Lect. Notes Phys.*, Springer (Berlin Heidelberg), 729, 111-150. D. Benest, C. Froeschlé, and E. Lega (Eds.), (2007)
- Wisdom, J. L., "The origin of the Kirkwood gaps: A mapping for asteroidal motion near the 3/1 commensurability.", *California Inst. of Tech., Pasadena*, (1981)